KING BRAND PIG STARTER-MEDICATED

ACTIVE DRUG INGREDIENT

INGREDIENTS

Ground Corn, Ground Wheat, Dehulled Soybean Meal, Distillers Dried Grains with Solubles, Wheat Millrun, Animal Fat, Calcium Carbonate, Dried Whey, Monocalcium & Dicalcium Phosphate, I-Lysine Hydrochloride, Salt, Choline Chloride, Zinc Sulfate, Ferrous Sulfate, Manganese Sulfate, Niacin Supplement, Vitamin E Supplement, Basic Copper Chloride, D-Calcium Pantothenate, Sodium Selenite, Menadione Dimethylpyrimidinol Bisulfite, Biotin Supplement, Riboflavin Supplement, Vitamin A Supplement, Mineral Oil, Vitamin D3 Supplement, Vitamin B12 Supplement, Ethylenediamine Dihydriodide, Folic Acid, Pyridoxine Hydrochloride, Thiamine Mononitrate, Hydrolyzed Yeast, L-Threonine, d I Methionine, Direct-Fed Microbials (Heat Stable cultures of Lactobacillus acidophilus, L. casei, Bifidobacterium bifidium, Enterococcus faecium and Aspergillus oryzae), Natural & Artificial Flavoring, Hydrated Calcium-Sodium Aluminosilicate, Copper Lysine Complex, Zinc Methionine Complex, Manganese Methionine Complex, Cobalt Glucoheptonate, Garlic Extract, Anise Extract, Horseradish, Juniper Extract, Chinese Cassia Bark Extract, Ginger Extract, Selenium Yeast,, Sweetener, Aspergillus niger Fermentation Product

FEEDING DIRECTIONS

1. Feed Pig Starter free choice as the sole ration to growing pigs 11 to 44 lbs bodyweight.

2. Store Pig Starter in a cool, dry area. Never feed moldy or insect-infested feed

WARNING: WITHDRAW FEED 2 DAYS BEFORE SLAUGHTER.

Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Lasalocid, Monensin, Narasin, Salinomycin, or Semduramycin) as adverse reactions may occur. If signs of toxicity occur, discontinue use. The effects of Tiamulin on swine reproductive performance, pregnancy, and lactation have not been determined. Use as sole source of tiamulin.

KING MEANS QUALITY

Manufactured By

L. A. HEARNE COMPANY- KING CITY, CA. 93930

H501

NET WEIGHT 22.6 kg (50lb)