

**UMBARGER-HEARNE  
BEEF BREEDER/GROWER SHOW RATION  
TYPE C MEDICATED**

DECOQUINATE.....22.7 Grams/Ton  
For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle

**NO WITHDRAWAL PERIOD BEFORE SLAUGHTER**

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	12.0%
Crude Fat, not less than.....	3.0%
Crude Fiber, not more than.....	10.0%
Calcium, not less than.....	0.5%
Calcium, not more than.....	1.0%
Phosphorus, not less than.....	0.35%
Salt, not less than.....	0.3%
Salt, not more than.....	0.8%
Potassium, not less than.....	0.65%
Vitamin A, not less than.....	.4000 IU/LB

**INGREDIENTS**

Steam Rolled Oats, Steam Rolled Corn, Cane Molasses & Vegetable Oil Blend, Steam Rolled Barley, Dehulled Soybean Meal, Cottonseed Hulls (not more than 5%), Linseed Meal, Calcium Carbonate, Cane Molasses, Salt, Wheat Millrun, Monocalcium & Dicalcium Phosphate, Corn Gluten Feed, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Cobalt Sulfate, Dried Aspergillus Oryzae Fermentation Product, Magnesium Oxide, Potassium Chloride, Ground Extruded Whole Soybeans, Zinc Proteinates, Monocalcium Phosphate, Dicalcium Phosphate, Vitamin D-3 Supplement, Vitamin E Supplement, Yeast Culture, Mineral Oil, Copper Proteinates, Choline Chloride, Vitamin A Supplement, Niacin Supplement, d-Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Thiamine Mononitrate, Natural and Artificial Flavor, Cobalt Carbonate., Mold Inhibitor (Propionic, Ammonium Hydroxide), Hydrolyzed Yeast, Processed Grain By-Products, Dried Lactobacillus Acidophilus Fermentation Product, Dried Lactobacillus Casei Fermentation Product, Dried Enterococcus Faecium Fermentation Product, Dried Bifidobacterium Longum Fermentation Product, Dried Aspergillus Oryzae Fermentation Extract, Fat Product Feed Grade and Hydrated Sodium Calcium Aluminosilicate, Potassium Magnesium Sulphate, Dried Vegetable Fat, Selenium Yeast, Vitamin D3 Supplement, Cobalt Carbonate, Condensed Lignin Sulfonate

**FEEDING DIRECTIONS**

Feed as a sole grain ration along with hay of choice. Free choice mineral and clean water should be made available at all times. Feed this Type C medicated feed at level to provide 22.7 mg Decoquinatone per 100 lb of bodyweight per day according to the table below.

<u>Steer Body Wt.</u>	<u>Daily Feeding Rate Lbs</u>	<u>Mg of Decoquinatone</u>
600 lbs	12.0 lb/day	136.2 mg
800 lbs	16.0 lb/day	181.6 mg
1000 lbs	20.0 lb/day	227 mg
1200 lbs	24.0 lb/day	272.4 mg

Store feed in a cool, dry area. Never feed moldy or insect-infested feed.

**WARNING:** Do not feed to cows producing milk for food. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed type C feed or milk replacer to provide 22.7 mg per 100 lb. bodyweight (0.5 mg/kg) per day.

**CAUTION:** Do not use in feeds containing bentonite.

**CAUTION:** Do not exceed 0.30 ppm added selenium from all supplemental sources in the total diet.

Manufactured By  
**L. A. HEARNE COMPANY- KING CITY, CA. 93930**  
(800) 253-7346 [www.king-brand.com](http://www.king-brand.com)  
**NET WEIGHT 22.6 kg (50 lb)**

120618

B192